

How to Describe Neutrino Mixing and CP Violation

Harald Fritzsch

Sektion Physik, Universität München, D-80333 München, Germany

Zhi-zhong Xing *

Institute of High Energy Physics, P.O. Box 918 (4), Beijing 100039, China

Abstract

We present a classification of possible parametrizations of the 3×3 lepton flavor mixing matrix in terms of the rotation and phase angles. A particular parametrization, which is most convenient to describe the observables of neutrino oscillations and that of the neutrinoless double beta decay, is studied in detail. The possibility to determine the Dirac- and Majorana-type CP-violating phases are also discussed.

*Electronic address: xingzz@mail.ihep.ac.cn

The accumulating evidence for atmospheric and solar neutrino oscillations strongly suggests that neutrinos are massive and lepton flavors are mixed [1]. In general the flavor mixing among N different lepton families is described by a $N \times N$ unitary matrix V , whose number of independent parameters depends on the nature of neutrinos. If neutrinos are Dirac particles, V can be parametrized in terms of $N(N-1)/2$ rotation angles and $(N-1)(N-2)/2$ phase angles. If neutrinos are Majorana particles, however, a full parametrization of V requires $N(N-1)/2$ rotation angles and the same number of phase angles. The flavor mixing of charged leptons and Dirac neutrinos is completely analogous to that of quarks, for which a number of different parametrizations have been proposed and classified in the literature [2]. One of the purposes of this paper is to classify the parametrizations of flavor mixing between charged leptons and Majorana neutrinos with $N = 3$. Regardless of the phase-assignment freedom, we find that there are nine distinct ways to describe the 3×3 lepton flavor mixing matrix.

Although different representations of lepton flavor mixing are mathematically equivalent, one of them is very likely to describe the underlying physics of lepton mass generation and CP violation in a more transparent way, or is particularly convenient in the analyses of experimental data on neutrino oscillations and lepton-number-violating processes. We point out that there does exist such a parametrization, which allows for simple connections between the measurable quantities and the flavor mixing angles. In addition, the phase assignment of this “standard” parametrization assures that the single Dirac-type CP-violating phase is associated only with neutrino oscillations, and the two Majorana-type CP-violating phases are associated only with the neutrinoless double beta decay. Some remarks are made on the difficulty to separately determine the Majorana-type CP-violating phases.

Let us start to parametrize the 3×3 lepton flavor mixing V , which is defined to link the neutrino flavor eigenstates $(\nu_e, \nu_\mu, \nu_\tau)$ to the neutrino mass eigenstates (ν_1, ν_2, ν_3) :

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \begin{pmatrix} V_{e1} & V_{e2} & V_{e3} \\ V_{\mu1} & V_{\mu2} & V_{\mu3} \\ V_{\tau1} & V_{\tau2} & V_{\tau3} \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}. \quad (1)$$

The strength of CP or T violation in normal neutrino oscillations, no matter whether neutrinos are Dirac or Majorana particles, depends only upon a universal parameter of V , defined as J through the following equation [3]:

$$\text{Im}(V_{\alpha i} V_{\beta j} V_{\alpha j}^* V_{\beta i}^*) = J \sum_{\gamma, k} \epsilon_{\alpha\beta\gamma} \epsilon_{ijk}, \quad (2)$$

where the Greek and Latin subscripts run over (e, μ, τ) for charged leptons and over $(1, 2, 3)$ for neutrinos, respectively.

In analogy to the quark mixing matrix, V can be expressed as a product of three unitary matrices O_1 , O_2 and O_3 , which correspond to simple rotations in the complex (1,2), (2,3) and (3,1) planes:

$$O_1(\theta_1, \alpha_1, \beta_1, \gamma_1) = \begin{pmatrix} c_1 e^{i\alpha_1} & s_1 e^{-i\beta_1} & 0 \\ -s_1 e^{i\beta_1} & c_1 e^{-i\alpha_1} & 0 \\ 0 & 0 & e^{i\gamma_1} \end{pmatrix},$$

$$\begin{aligned}
O_2(\theta_2, \alpha_2, \beta_2, \gamma_2) &= \begin{pmatrix} e^{i\gamma_2} & 0 & 0 \\ 0 & c_2 e^{i\alpha_2} & s_2 e^{-i\beta_2} \\ 0 & -s_2 e^{i\beta_2} & c_2 e^{-i\alpha_2} \end{pmatrix}, \\
O_3(\theta_3, \alpha_3, \beta_3, \gamma_3) &= \begin{pmatrix} c_3 e^{i\alpha_3} & 0 & s_3 e^{-i\beta_3} \\ 0 & e^{i\gamma_3} & 0 \\ -s_3 e^{i\beta_3} & 0 & c_3 e^{-i\alpha_3} \end{pmatrix},
\end{aligned} \tag{3}$$

where $s_i \equiv \sin \theta_i$ and $c_i \equiv \cos \theta_i$ (for $i = 1, 2, 3$). Obviously $O_i O_i^\dagger = O_i^\dagger O_i = 1$ holds, and any two rotation matrices do not commute with each other. Note that the matrix $O_i^\dagger$ or O_i^{-1} plays an equivalent role as O_i in constructing V , because of

$$O_i^\dagger(\theta_i, \alpha_i, \beta_i, \gamma_i) = O_i^{-1}(\theta_i, \alpha_i, \beta_i, \gamma_i) = O_i(-\theta_i, -\alpha_i, -\beta_i, -\gamma_i). \tag{4}$$

Note also that

$$O_i(\theta_i, \alpha_i, \beta_i, \gamma_i) \otimes O_i(\theta'_i, \alpha'_i, \beta'_i, \gamma'_i) = O_i(\theta''_i, \alpha''_i, \beta''_i, \gamma''_i) \tag{5}$$

holds, where $\theta''_i, \alpha''_i, \beta''_i$ and γ''_i are simple functions of $(\theta_i, \alpha_i, \beta_i, \gamma_i)$ and $(\theta'_i, \alpha'_i, \beta'_i, \gamma'_i)$. In particular, one will get $\theta''_i = \theta_i + \theta'_i$ if all the complex phases in O_i are switched off. Thus the product $O_i(\theta_i) \otimes O_i(\theta'_i) \otimes O_j(\theta_j)$ or $O_i(\theta_i) \otimes O_j(\theta_j) \otimes O_j(\theta'_j)$ is unable to cover the whole space of a 3×3 unitary matrix and should be excluded. We find that there are only twelve different possibilities to arrange the product of O_1, O_2 and O_3 , which can cover the whole 3×3 space and provide a full description of V . Explicitly, six of the twelve different combinations of O_i belong to the category

$$V = O_i(\theta_i, \alpha_i, \beta_i, \gamma_i) \otimes O_j(\theta_j, \alpha_j, \beta_j, \gamma_j) \otimes O_i(\theta'_i, \alpha'_i, \beta'_i, \gamma'_i) \tag{6}$$

with $i \neq j$, where the complex rotation matrix O_i occurs twice; and the other six belong to the category

$$V = O_i(\theta_i, \alpha_i, \beta_i, \gamma_i) \otimes O_j(\theta_j, \alpha_j, \beta_j, \gamma_j) \otimes O_k(\theta_k, \alpha_k, \beta_k, \gamma_k) \tag{7}$$

with $i \neq j \neq k$, in which the rotations take place in three different complex planes.

It should be noted that only nine of the twelve parametrizations, three from Eq. (6) and six from Eq. (7), are structurally different. The reason is simply that the combinations $O_i \otimes O_j \otimes O_i$ and $O_i \otimes O_k \otimes O_i$ (for $i \neq k$) in Eq. (6) are correlated with each other [4]. To see this point clearly, we switch off the relevant phase parameters in O_i and then arrive at the following relations:

$$\begin{aligned}
O_1(\theta_1) \otimes O_3(\theta_3) \otimes O_1(\theta'_1) &= O_1(\theta_1 + 90^\circ) \otimes O_2(\theta_2 = \theta_3) \otimes O_1(\theta'_1 - 90^\circ), \\
O_2(\theta_2) \otimes O_1(\theta_1) \otimes O_2(\theta'_2) &= O_2(\theta_2 + 90^\circ) \otimes O_3(\theta_3 = \theta_1) \otimes O_2(\theta'_2 - 90^\circ), \\
O_3(\theta_3) \otimes O_2(\theta_2) \otimes O_3(\theta'_3) &= O_3(\theta_3 + 90^\circ) \otimes O_1(\theta_1 = \theta_2) \otimes O_3(\theta'_3 - 90^\circ).
\end{aligned} \tag{8}$$

Therefore three of the six combinations in Eq. (6) need not be treated as independent choices, even though the phase parameters are taken into account. We then conclude that there exist nine distinct parametrizations of the 3×3 lepton flavor mixing matrix V , no matter how the complex phases are arranged among the nine elements of V .

In each of the nine distinct parametrizations for V , there apparently exist nine phase parameters. Six of them or combinations thereof can be absorbed by redefining the arbitrary phases of charged lepton fields and a common phase of neutrino fields. If neutrinos are Dirac particles, one can also redefine the arbitrary phases of Dirac neutrino fields to reduce the number of the remaining phase parameters from three to one. In this case V consists of only a single nontrivial phase parameter, which violates CP symmetry. If neutrinos are Majorana particles, however, there is no freedom to rearrange the relative phases of Majorana neutrino fields. Hence V is totally composed of three nontrivial phase parameters in the latter case. There is much freedom, through redefinition of the arbitrary phases of charged lepton fields, to place the three CP-violating phases among the nine elements of V . In particular, it is always possible to parametrize the Majorana-type flavor mixing matrix as a product of the Dirac-type flavor mixing matrix (with three mixing angles and a single CP-violating phase) and a diagonal phase matrix (with two unremovable CP-violating phases) [5].

To be more specific, let us take two typical examples to show the parametrization of V in terms of three mixing angles and three CP-violating phases.

Example A: The lepton flavor mixing matrix can be parametrized, in close analogy to a representation of the quark mixing matrix [6], as follows:

$$V = \begin{pmatrix} s_l s_\nu c + c_l c_\nu e^{-i\phi} & s_l c_\nu c - c_l s_\nu e^{-i\phi} & s_l s \\ c_l s_\nu c - s_l c_\nu e^{-i\phi} & c_l c_\nu c + s_l s_\nu e^{-i\phi} & c_l s \\ -s_\nu s & -c_\nu s & c \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{i\rho} & 0 \\ 0 & 0 & e^{i\sigma} \end{pmatrix}, \quad (9)$$

where $s_l \equiv \sin \theta_l$, $c_\nu \equiv \cos \theta_\nu$, etc. The three mixing angles $(\theta_l, \theta_\nu, \theta)$ may have simple physical interpretations in a specific scheme of lepton mass matrices [7]. In particular, we expect θ_l to be small in magnitude, as a natural consequence of the mass hierarchy of three charged leptons. It is obvious that only the phase ϕ remains present, if neutrinos are assumed to be Dirac particles. The reason is simply that the diagonal phase matrix on the right-hand side of Eq. (9), which consists of the Majorana-type CP-violating phases ρ and σ , can be rotated away by redefining the phases of Dirac neutrino fields. In other words, only ϕ is associated with CP or T violation in normal neutrino oscillations (measured by $J = s_l c_l s_\nu c_\nu s^2 c \sin \phi$), no matter whether neutrinos are Majorana particles or not. The diagonal phase matrix of V signifies the Majorana nature of neutrinos and affects the neutrinoless double beta decay and some other lepton-number-violating processes.

Example B: The lepton flavor mixing matrix can be parametrized, in a form similar to the parametrization of quark flavor mixing discussed in Ref. [8], as follows:

$$V = \begin{pmatrix} c_1 c_3 & s_1 c_3 & s_3 \\ -c_1 s_2 s_3 - s_1 c_2 e^{-i\delta} & -s_1 s_2 s_3 + c_1 c_2 e^{-i\delta} & s_2 c_3 \\ -c_1 c_2 s_3 + s_1 s_2 e^{-i\delta} & -s_1 c_2 s_3 - c_1 s_2 e^{-i\delta} & c_2 c_3 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{i\rho} & 0 \\ 0 & 0 & e^{i\sigma} \end{pmatrix} \quad (10)$$

with $s_i \equiv \sin \theta_i$ and $c_i \equiv \cos \theta_i$ (for $i = 1, 2, 3$). Note that the location of the Dirac-type CP-violating phase δ in V is different from that advocated by the Particle Data Group [9]. The advantage of our present phase assignment is that δ itself does not appear in the effective Majorana mass term of the neutrinoless double beta decay, as one can see later on. Without loss of generality, the three mixing angles $(\theta_1, \theta_2, \theta_3)$ can all be arranged to lie in the first

quadrant. Arbitrary values between -180° and $+180^\circ$ are allowed for δ and the Majorana-type CP-violating phases ρ and σ . The CP- and T-violating effects in normal neutrino oscillations are measured by $J = s_1 c_1 s_2 c_2 s_3 c_3^2 \sin \delta$. As the magnitude of J is independent of the specific parametrizations of V , one can easily find out the relation between the Dirac-type CP-violating phases ϕ in Eq. (8) and δ in Eq. (10): $\sin \delta / \sin \phi = (s_l c_l s_\nu c_\nu s^2 c) / (s_1 c_1 s_2 c_2 s_3 c_3^2)$.

Of course, both examples taken above and other possible parametrizations of V are mathematically equivalent, and adopting any of them does not have any specific physical significance. It is quite likely, however, that one particular parametrization is more useful and transparent than the others in the analyses of data from various neutrino experiments and (or) towards a deeper understanding of the underlying dynamics responsible for lepton mass generation and CP violation.

We find that Example B is very convenient to confront with the observables of neutrino oscillations and that of the neutrinoless double beta decay (see the next section for a detailed discussion). In particular, it is favored if the solar neutrino problem invokes a large-angle MSW (or vacuum oscillation) solution [10]. In this case, the lepton mixing matrix V is expected to be roughly symmetric about its axis V_{e3} - $V_{\mu 2}$ - $V_{\tau 1}$; i.e., $\theta_1 \sim \theta_2$ holds. If the small-angle MSW oscillation were the true solution to the solar neutrino problem[†], however, Example A would show its advantages. For instance, V would be expected to be roughly symmetric about its axis V_{e1} - $V_{\mu 2}$ - $V_{\tau 3}$ in the latter case (i.e., $\theta_l \sim \theta_\nu$ holds), just like the approximate off-diagonal symmetry of the 3×3 quark flavor mixing matrix about its axis V_{ud} - V_{cs} - V_{tb} [13]. Both θ_l and θ_ν might get simple physical interpretations in terms of the ratios of charged lepton and neutrino masses, provided that the texture of lepton mass matrices is constrained by some flavor symmetries. An instructive possibility is

$$\tan \theta_l \approx \sqrt{\frac{m_e}{m_\mu}} \sim \mathcal{O}(10^{-2}), \quad \tan \theta_\nu \approx \sqrt{\frac{m_1}{m_2}} \sim \mathcal{O}(10^{-2}), \quad (11)$$

if the neutrino masses exhibit a similar hierarchy as the charged lepton masses or the quark masses.

Furthermore, it is worth pointing out a useful relation in the limit $|V_{e3}| = 0$ (i.e., $\theta_3 = 0$ or $\theta_l = 0$):

$$\left| \frac{V_{e2}}{V_{e1}} \right| = \left| \frac{V_{\mu 1}}{V_{\mu 2}} \right| = \left| \frac{V_{\tau 1}}{V_{\tau 2}} \right| = \begin{cases} \tan \theta_\nu & \text{(Example A)} \\ \tan \theta_1 & \text{(Example B)} \end{cases}. \quad (12)$$

Such a result is meaningful, because small $|V_{e3}|$ is favored by current experimental data on neutrino oscillations.

Let us concentrate on the parametrization in Eq. (10) (i.e., Example B) and confront it with the measurable quantities of lepton flavor mixing and CP violation. First of all,

[†]The latest SNO experiment [11], together with the Super-Kamiokande data, has provided the first direct evidence that there is a muon- and (or) tau-neutrino component in the solar electron-neutrino flux. The global fit shows that the large-angle MSW solution is apparently favored and the small-angle MSW solution is highly disfavored [12]. However, it remains too early to claim that the small-angle MSW solution has been convincingly ruled out. Further experimental effort is desirable to pin down the true solution to the solar neutrino puzzle.

the mixing angle θ_3 can be determined from measuring the survival probability of electron neutrinos at the scale of atmospheric neutrino oscillations in a long-baseline (LBL) neutrino experiment:

$$\sin^2 2\theta_{\text{LBL}} \approx 4|V_{e3}|^2 (1 - |V_{e3}|^2) = 4s_3^2 (1 - s_3^2) = \sin^2 2\theta_3. \quad (13)$$

The current constraint obtained from CHOOZ [14] and Palo Verde [15] reactor experiments, $\sin^2 2\theta_{\text{LBL}} \ll 1$, indicates that θ_3 may be quite small. This result, together with the mass-squared hierarchy $\Delta m_{\text{sun}}^2 \ll \Delta m_{\text{atm}}^2$ showing up in a variety of analyses of the experimental data on atmospheric and solar neutrino oscillations [1, 10], strongly implies that solar and atmospheric neutrino oscillation phenomena approximately decouple from each other. Hence the mixing angles θ_1 and θ_2 essentially measure the corresponding amplitudes of solar ($\nu_e \rightarrow \nu_e$) and atmospheric ($\nu_\mu \rightarrow \nu_\mu$) neutrino oscillations; i.e.,

$$\begin{aligned} \sin^2 2\theta_{\text{sun}} &\approx 4|V_{e1}|^2 |V_{e2}|^2 = 4s_1^2 c_1^2 c_3^4 \approx \sin^2 2\theta_1, \\ \sin^2 2\theta_{\text{atm}} &\approx 4|V_{\mu 3}|^2 (1 - |V_{\mu 3}|^2) = 4s_2^2 c_3^2 (1 - s_2^2 c_3^2) \approx \sin^2 2\theta_2. \end{aligned} \quad (14)$$

We see that all three mixing angles of the parametrization in Eq. (10) have simple relations to measurable quantities ($\theta_1 \approx \theta_{\text{sun}}$, $\theta_2 \approx \theta_{\text{atm}}$, and $\theta_3 \approx \theta_{\text{LBL}}$), at least in the leading-order approximation.

The Dirac-type CP-violating phase δ can be determined from CP- and (or) T-violating asymmetries in normal long-baseline neutrino oscillations. In vacuum, the T-violating asymmetry between the probabilities of $\nu_\alpha \rightarrow \nu_\beta$ and $\nu_\beta \rightarrow \nu_\alpha$ transitions amounts to the CP-violating asymmetry between the probabilities of $\nu_\alpha \rightarrow \nu_\beta$ and $\bar{\nu}_\alpha \rightarrow \bar{\nu}_\beta$ transitions [16]:

$$\begin{aligned} \Delta P &\equiv P(\nu_\alpha \rightarrow \nu_\beta) - P(\bar{\nu}_\alpha \rightarrow \bar{\nu}_\beta) \\ &= P(\nu_\alpha \rightarrow \nu_\beta) - P(\nu_\beta \rightarrow \nu_\alpha) \\ &= -16J \sin F_{21} \cdot \sin F_{31} \cdot \sin F_{32}, \end{aligned} \quad (15)$$

where the subscripts (α, β) run over (e, μ) , (μ, τ) or (τ, e) , and $F_{ij} \equiv 1.27 \Delta m_{ij}^2 L / E$ with $\Delta m_{ij}^2 \equiv m_i^2 - m_j^2$ being the mass-squared differences of neutrinos (in unit of eV^2), L being the baseline length (in unit of km), and E being the neutrino beam energy (in unit of GeV). A determination of J from ΔP will allow us to extract the CP-violating phase δ , provided that all three mixing angles $(\theta_1, \theta_2, \theta_3)$ have been measured elsewhere. In practice, however, all these measurable quantities may be contaminated due to the presence of terrestrial matter effects. Hence the fundamental parameters of lepton flavor mixing need be disentangled from the matter-corrected ones.

Regardless of the Majorana-type phases ρ and σ , which have nothing to do with normal neutrino oscillations, we have located the Dirac-type phase δ in such a way that the matrix elements in the first row and the third column of V are real. As a consequence, the CP-violating phase δ does not appear in the effective mass term of the neutrinoless double beta decay. Indeed the latter reads:

$$\begin{aligned} \langle m \rangle_{ee} &= |m_1 V_{e1}^2 + m_2 V_{e2}^2 + m_3 V_{e3}^2| \\ &= \sqrt{\mathbf{a} + \mathbf{b} \cos 2\rho + \mathbf{c} \cos 2\sigma + \mathbf{d} \cos 2(\rho - \sigma)}, \end{aligned} \quad (16)$$

where

$$\begin{aligned}
\mathbf{a} &= m_1^2 c_1^4 c_3^4 + m_2^2 s_1^4 c_3^4 + m_3^2 s_3^4, \\
\mathbf{b} &= 2m_1 m_2 s_1^2 c_1^2 c_3^4, \\
\mathbf{c} &= 2m_1 m_3 c_1^2 s_3^2 c_3^2, \\
\mathbf{d} &= 2m_2 m_3 s_1^2 s_3^2 c_3^2.
\end{aligned} \tag{17}$$

It becomes obvious that $\langle m \rangle_{ee}$ is independent of both the mixing angle θ_2 and the CP-violating phase δ . On the other hand, CP- and T-violating asymmetries in normal neutrino oscillations depend only upon the Dirac-type phase δ or the universal CP-violating parameter J ; i.e., they have nothing to do with the Majorana-type CP-violating phases ρ and σ [17]. We then arrive at the conclusion that the two different types of CP-violating phases can (in principle) be studied in two different types of experiments.

It is worth pointing out that the expression of $\langle m \rangle_{ee}$ can particularly be simplified, if $\theta_3 \approx 0$ and $m_1 \approx m_2 \approx m_3$ are assumed. In this special case, we arrive at

$$\langle m \rangle_{ee} \approx m_1 \sqrt{1 - \sin^2 2\theta_1 \sin^2 \rho}, \tag{18}$$

which depends only upon a single Majorana phase ρ .

A long-standing and important question is whether the two Majorana phases ρ and σ can be separately determined by measuring other possible lepton-number-nonconserving processes, in addition to the neutrinoless double beta decay. While the answer to this question is affirmative in principle, it seems to be negative in practice. The key problem is that those lepton-number-violating processes, in which the Majorana phases can show up, are suppressed in magnitude by an extremely small factor compared to normal weak interactions. Therefore it is extremely difficult, even impossible, to measure or constrain ρ and σ in any experiment other than the one associated with the neutrinoless double beta decay.

To illustrate the above-mentioned difficulty in measuring ρ and σ separately, let us take a ‘‘Gedanken’’ experiment of neutrino-antineutrino oscillations for example [18]. We suppose a beam of positive muons to be incident upon a neutron target, from which the antineutrinos $\bar{\nu}_\mu$ are emitted through the usual W -mediated reaction $\mu^+ + n \rightarrow \bar{\nu}_\mu + p$ in a given direction. Such an energetic beam of antineutrinos ($E \gg m_i$ for $i = 1, 2, 3$) is then arranged to hit another neutron target at proper time t , leading to the emission of negative muons through the lepton-number-conserving reaction $\nu_\mu + n \rightarrow \mu^- + p$. The overall process can actually happen, because of the lepton-number-violating conversion of $\bar{\nu}_\mu$ into ν_μ at the interval t . The effective amplitude of this $\bar{\nu}_\mu \rightarrow \nu_\mu$ oscillation is therefore expressed as

$$A(\bar{\nu}_\mu \rightarrow \nu_\mu) = \frac{1}{E} \sum_{k=1}^3 \left[m_k (V_{\mu k}^*)^2 e^{-iE_k t} \right], \tag{19}$$

where E is the neutrino beam energy and $V_{\mu k}$ (for $k = 1, 2, 3$) are the elements of the flavor mixing matrix V . Obviously $|A(\bar{\nu}_\mu \rightarrow \nu_\mu)|^2$ depends upon all of the three mixing angles and the three CP-violating phases in the parametrization advocated above. It is therefore possible, in principle, to determine the Majorana phases ρ and σ separately from Eqs. (16)

and (19), if the Dirac-type phase (δ) and three mixing angles ($\theta_1, \theta_2, \theta_3$) have been fixed from the experiments of normal neutrino-neutrino and antineutrino-antineutrino oscillations. The probability of observing the $\bar{\nu}_\mu \rightarrow \nu_\mu$ oscillation is nevertheless suppressed by the factors $(m_k/E)^2$ in comparison with that of observing the normal $\nu_\mu \rightarrow \nu_\mu$ or $\bar{\nu}_\mu \rightarrow \bar{\nu}_\mu$ oscillation, whose amplitude is associated only with $|V_{\mu k}|^2$. As the factors m_k/E (for $k = 1, 2, 3$) are expected to be extremely tiny (e.g., of order 10^{-9} for $m_k \sim 1$ eV and $E \sim 1$ GeV), it is practically impossible to measure the $\bar{\nu}_\mu \rightarrow \nu_\mu$ oscillation. It is also extremely difficult or impossible to observe other similar types of neutrino-antineutrino oscillating effects [19].

It seems that one would have no way to determine all of the three CP-violating phases of V , even though the Majorana nature of neutrinos could finally be established (e.g., from the experiment on the neutrinoless double beta decay). Whether this will be the case has to be seen. On the theoretical side, how to predict or calculate those flavor mixing angles and CP-violating phases on a solid dynamical ground remains an open question. Phenomenologically, the parametrization in Eq. (10) is expected to be very useful and convenient, and might even be able to provide some insight into the underlying physics of lepton mass generation. We therefore recommend it to experimentalists and theorists as a standard parametrization of the 3×3 lepton flavor mixing matrix.

In summary, we have classified possible parametrizations of the 3×3 lepton flavor mixing matrix in terms of the rotation and phase angles. A particular parametrization, which is most convenient to confront with the measurables of neutrino oscillations and that of the neutrinoless double beta decay, has been emphasized from the phenomenological point of view.

Although the present non-accelerator neutrino experiments have yielded some impressive constraints on three lepton flavor mixing angles (θ_1 , θ_2 , and θ_3), a precise determination of them and a measurement of the Dirac-type CP-violating phase δ have to rely on a new generation of accelerator experiments with very long baselines, including the possible neutrino factories. In such long- or medium-baseline neutrino experiments the terrestrial matter effects, which may deform the oscillating behaviors of neutrinos in vacuum and even fake the genuine CP-violating signals, must be taken into account [20].

We expect that some significant progress can be made in our understanding of the lepton mass generation, flavor mixing and CP violation, once the precision measurements of neutrino oscillations are carried out in the long-baseline neutrino experiments. Nevertheless, it seems essentially impossible to separately determine the two Majorana-type CP-violating phases from any feasible measurements of the lepton-number-violating processes.

References

- [1] Super-Kamiokande Collaboration, Y. Fukuda *et al.*, Phys. Rev. Lett. **81** (1998) 1562; **81** (1998) 4279; <http://www-sk.icrr.u-tokyo.ac.jp/dpc/sk/>.
- [2] For a review, see: H. Fritzsch and Z.Z. Xing, Prog. Part. Nucl. Phys. **45** (2000) 1; hep-ph/9912358.

- [3] C. Jarlskog, Phys. Rev. Lett. **55** (1985) 1039.
- [4] H. Fritzsch and Z.Z. Xing, Phys. Rev. D **57** (1998) 594.
- [5] J. Schechter and J.W.F. Valle, Phys. Rev. D **22** (1980) 2227; S.M. Bilenky, J. Hosek, and S.T. Petcov, Phys. Lett. B **94** (1980) 495.
- [6] H. Fritzsch and Z.Z. Xing, Phys. Lett. B **413** (1997) 396.
- [7] See, e.g., H. Fritzsch and Z.Z. Xing, Phys. Lett. B **372** (1996) 265; Phys. Lett. B **440** (1998) 313; Phys. Rev. D **61** (2000) 073016; Z.Z. Xing, Phys. Rev. D **63** (2001) 057301.
- [8] L. Maiani, in *Proceedings of International Symposium on Lepton and Photon Interactions at High Energies* (DESY, Hamburg, 1977), p. 867; L.L. Chau and W.Y. Keung, Phys. Rev. Lett. **53** (1984) 1802; H. Fritzsch, Phys. Rev. D **32** (1985) 3058; H. Harari and M. Leurer, Phys. Lett. B **181** (1986) 123.
- [9] Particle Data Group, D.E. Groom *et al.*, Eur. Phys. J. C **15** (2000) 1.
- [10] Y. Takeuchi and T. Toshito, plenary talks given at ICHEP2000, Osaka, July 2000; <http://ichep2000.hep.sci.osaka-u.ac.jp/>.
- [11] SNO Collaboration, Q.R. Ahmad *et al.*, nucl-ex/0106015.
- [12] V. Barger, D. Marfatia, and K. Whisnant, hep-ph/0106207; G.L. Fogli, E. Lisi, D. Montanino, and A. Palazzo, hep-ph/0106247; J.N. Bahcall, M.C. Gonzalez-Garcia, and C. Peña-Garay, hep-ph/0106258.
- [13] H. Fritzsch and Z.Z. Xing, hep-ph/9903499; Z.Z. Xing, Nucl. Phys. B (Proc. Suppl.) **85** (2000) 187.
- [14] CHOOZ Collaboration, M. Apollonio *et al.*, Phys. Lett. B **420** (1998) 397; Phys. Lett. B **466** (1999) 415.
- [15] Palo Verde Collaboration, F. Boehm *et al.*, Phys. Rev. Lett. **84** (2000) 3764; Phys. Rev. D **62** (2000) 072002.
- [16] N. Cabibbo, Phys. Lett. B **72** (1978) 333; V. Barger, K. Whisnant, and R.J.N. Phillips, Phys. Rev. Lett. **45** (1980) 2084.
- [17] B. Kayser, hep-ph/0010065; S.M. Bilenky, S. Pascoli, and S.T. Petcov, hep-ph/0102265.
- [18] J. Schechter and J.W.F. Valle, Phys. Rev. D **23** (1981) 1666.
- [19] M. Flanz, W. Rodejohann, and K. Zuber, Eur. Phys. J. C **16** (2000) 453; and references therein.
- [20] See, e.g., B. Autin *et al.*, CERN-SPSC-98-30 (1998); M.G. Catanesi *et al.*, CERN-SPSC-99-35 (1999); D. Ayres *et al.*, physics/9911009; C. Albright *et al.*, hep-ex/0008064.